

Homework #6 Solutions

Monday, May 24, 2010
2:59 PM

11.9 \vec{P} is along \vec{y} (symmetry)

$$dP_y = 2b^2 \cdot \lambda_0 \sin^3 \varphi \cdot d\varphi$$

$$P_y = 2b^2 \lambda_0 \cdot \int_0^{\pi} \sin^3 \varphi \cdot d\varphi =$$

$$= 2b^2 \lambda_0 \cdot \frac{\pi}{2} = \pi b^2 \lambda_0$$

$$\vec{P}(t) = \pi b^2 \lambda_0 (\sin \omega t \cdot \hat{x} + \cos \omega t \cdot \hat{y})$$

$$\vec{\dot{P}}(t) = -\omega^2 \pi b^2 \lambda_0 (\sin \omega t \hat{x} + \cos \omega t \hat{y})$$

$$\langle |\dot{P}(t)|^2 \rangle = \omega^4 \pi^2 b^4 \lambda_0^2$$

$$\text{Power} = \frac{\mu_0}{6\pi c} \cdot (\omega^2 \pi b^2 \lambda_0)^2 = \frac{\mu_0 \pi \omega^4 b^4 \lambda_0^2}{6c}$$

11.10

$$U = m \cdot g \cdot h \quad t = \sqrt{\frac{2h}{g}}$$

$$\text{Power} = \frac{\mu_0 g^2 a^2}{6\pi c}$$

$$\text{Energy dissipated} = \text{Power} \cdot t =$$

$$= \frac{\mu_0 q^2 \cdot g^2}{6\pi c} \cdot \sqrt{\frac{2L}{g}}$$

$$\frac{\text{Energy dissip.}}{U} = \frac{\mu_0 q^2}{6\pi c \cdot mgh} \cdot \sqrt{\frac{2h}{g}} =$$

$$= \frac{\mu_0 q^2}{6\pi c \cdot m} \cdot \sqrt{\frac{2g}{h}} \approx 10^{-24}$$

II.13 $P_{\text{Power}} = \frac{\mu_0 q^2 a^2}{6\pi c}$

$$t = \frac{V_o}{a} \Rightarrow E = P_{\text{Power}} \cdot t = \frac{\mu_0 q^2 \cdot a \cdot V_o}{6\pi c}$$

$$\frac{E}{\frac{mv^2}{2}} = \frac{\mu_0 q^2 \cdot a}{3\pi m V_o c}$$

b) $\frac{E}{\frac{mv^2}{2}} = 1.3 \cdot 10^{10} \text{ (small)}$

II.25

| | | |
|--------|------|---|
| • q | • -q | $P = q \cdot 2z$ $\dot{P} = q \cdot 2 \dot{z}$ |
|--------|------|---|

$$P = \frac{\mu_0 (\ddot{P})^2}{6\pi c} \quad m \ddot{z} = \frac{q^2}{4\pi \epsilon_0 (zz)^2}$$

$$P = \frac{\mu_0}{6\pi c} \cdot \left(\frac{q^3}{8\pi\epsilon_0 m \tau^2} \right)^2 =$$
$$= \left(\frac{\mu_0 c q^2}{4\pi} \right)^2 \cdot \frac{1}{6m^2 \epsilon^4}$$